





CLINICAL SCIENCE

Factors predicting axial spondyloarthritis among first-degree relatives of probands with ankylosing spondylitis: a family study spanning 35 years

Sjef M van der Linden ^{1,2}, Muhammad Asim Khan ³, Zhixiu Li ^{4,5}, Heinz Baumberger,⁶ Hermine van Zandwijk,⁷ Kazim Khan,⁸ Peter M Villiger ⁹, Matthew A Brown ^{10,11}

Handling editor Josef S Smolen

For numbered affiliations see end of article.

Correspondence to

Professor Sjef M van der Linden, University of Bern, Bern, Switzerland; svdlinden@scarlet.be
Professor Matthew A Brown; matt.brown@genomicsengland.co.uk

HB Deceased

Received 27 December 2021
Accepted 22 February 2022

ABSTRACT

Objective Factors predicting axial spondyloarthritis (axSpA) among first-degree relatives (FDRs) of ankylosing spondylitis (AS) patients need to be defined. We investigated the predictive value of the probands' HLA-B27 and radiographic sacroiliitis status on disease occurrence among their FDR. We also assessed the predictive value of features of the clinical history, including chronic inflammatory back pain (CIBP) and acute anterior uveitis (AAU), among the FDR and how they can be used to improve classification and diagnosis of axSpA.

Methods In 1985, we studied 363 AS probands and 806 FDR who underwent rheumatologic examination, completed questionnaires, provided blood samples for HLA-typing and underwent radiography of sacroiliac joints. At follow-up in 2018–2019, 125 patients and 360 FDR were available for study, and completed a postal questionnaire about axSpA features. FDRs were asked to report whether after 1985 they had been diagnosed by Swiss rheumatologists as having axSpA.

Results Among HLA-B27(+) FDR, axSpA occurred in 25.4%–26.3%, independent of the radiographic sacroiliitis status of the proband. AAU occurred in 13/34 (38.2%) FDR with axSpA vs 29/251 (11.6%) FDR without axSpA ($p=0.00004$, $OR=4.74$ 95% CI 2.15 to 10.47). The presence of CIBP at baseline did not predict later occurrence of axSpA but combining CIBP and pain/discomfort at the thoracic spine *and* at anterior (ventral) chest wall ever, assessed at follow-up in 2018–2019, provided 83.1% sensitivity and 87.2% specificity for current axSpA.

Conclusion Occurrence of AAU among FDR of axSpA probands should prompt screening for axSpA. Moreover, co-occurrence of CIBP and pain/discomfort in the thoracic spine and at anterior chest wall as a three-question tool may further enhance clinical suspicion of axSpA among these FDR.

INTRODUCTION

The high familiarity of ankylosing spondylitis (AS) is well established.^{1–3} The prevalence of AS is considerably higher among HLA-B27(+) first-degree relatives (FDR) of patients with the disease compared with HLA-B27(+) persons from the general population.^{4,5} This points to an important role of additional MHC genes, such as HLA-B60, and many

Key messages

What is already known about this subject?

- Familial occurrence of ankylosing spondylitis is well known, but factors predicting axial spondyloarthritis (axSpA) occurrence among first-degree relatives (FDR) are less well identified.

What does this study add?

- Occurrence of chronic inflammatory back pain (CIBP) in FDR of axSpA probands is not a reliable predictor of later development of axSpA. However, occurrence of acute anterior uveitis and/or a combined occurrence of CIBP and pain/discomfort in the thoracic spine and at anterior chest wall, enhance early clinical suspicion and early diagnosis of axSpA among FDR of axSpA patients.
- 'Healthy' HLA-B27(+) FDR may have unnoticed 'hidden' axSpA, in particular if they have thoracic complaints in the absence of CIBP.

How might this impact on clinical practice or future developments?

- A three-question tool combining pain and discomfort at lumbar spine, thoracic spine and anterior chest wall has high sensitivity (83.1%) and specificity (87.2%). However, this triad needs to be fully validated as a potential diagnostic tool.

non-MHC genes in contributing to disease susceptibility.^{6,7}

Nowadays, the notion of AS has changed with increased emphasis on early diagnosis at its non-radiographic phase.^{8,9} This has led to the concept of axial spondyloarthritis (axSpA), comprising both radiographic axSpA (AS defined by the modified New York (mNY) criteria¹⁰ and non-radiographic axSpA (nr-axSpA). Whether the recurrence risk in relatives of axSpA patients varies according to the radiographic status of the proband is unknown. With nr-axSpA reported to be 2–3 times¹¹ more prevalent than AS itself, there is a significant clinical need to determine this risk rate, not the least for counselling and screening purposes.



© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: van der Linden SM, Khan MA, Li Z, *et al.* *Ann Rheum Dis* Epub ahead of print: [please include Day Month Year]. doi:10.1136/annrheumdis-2021-222083

We have recently completed a family study spanning 35 years, the longest longitudinal study in axSpA to date. Here, we report features from patients' clinical history that best predict presence of axSpA for the large number of FDR who developed the disease during these 35 years. We evaluated whether presence at early age of chronic inflammatory low back pain (CIBP) and occurrence of features of acute anterior uveitis (AAU) are associated with onset of axSpA.

METHODS

Here the term 'axSpA' comprises the full spectrum of axSpA, that is, both radiographic (AS defined by the mNY criteria) and nr-axSpA.¹⁰

First phase of the study

In 1985, all members of the nation-wide Swiss Ankylosing Spondylitis Patient Society were invited to participate in the family study together with their spouses and FDR. These relatives were invited irrespective of whether they were known to have any rheumatic disease. The study was performed in centres spread all over the country.

A total of 1178 persons consented to participate and completed questionnaires on disease manifestations. They also underwent physical examination of axial and peripheral joints by a rheumatologist. Peripheral blood nucleated cells (PBNCs) were stored for HLA-typing and subsequent genetic analysis. Furthermore, to assess the presence of sacroiliitis, consenting non-pregnant participants, aged 18 and over, underwent pelvic radiography unless a recent radiograph was available. Pelvic radiographs were available for 360 of the 363 probands; the three probands with missing pelvic radiographs were excluded in this analysis.

Each sacroiliac (SI) joint on pelvic radiographs (total number 1081) was 'blindly' assessed twice by each of 4 experienced readers, that is, 8 (occasionally 9) times. The radiographs of 163/360 probands and 22/713 FDR were only available on-site for a few hours at the time of participant's physical examination in the local hospital, and therefore, could only be assessed once. Overall, 17.2% of the 1081 radiographs were read once, 0.4% 2–4 times, 3.2% 5–7 times and 79.2% 8–9 times. The sacroiliitis score ranged from 0 (normal) to 4 (ankylosis) for each SI joint assessment by a reader as per the mNY scoring system.¹⁰ All scores for a single SI joint were added and divided by the number of assessments (range 1–9). Scores below bilateral grade 2.0 or unilateral grade 3.0 were considered not fulfilling the mNY criteria. Interobserver and intraobserver reliability were assessed by evaluating a subset of 243 pelvic films. Observers read films twice in sets of 40–50 radiographs. The interval between both readings was ≥ 7 days. The interobserver and intraobserver reliability coefficients were 0.865 and 0.903, respectively.

In 1986, all participants were informed by a letter about their individual HLA-B27 test result and radiographic findings of the SI joints.

Follow-up study

The former participants were asked to complete a 157-item postal questionnaire on manifestations of AS. The questionnaire dealt particularly with current or past symptoms at lumbar and gluteal region, thoracic spine and front part of the chest, and also addressed symptoms suggesting episodes of AAU. The next decisive step to perform the follow-up study was to determine whether the former 1178 participants were still alive and, if so, to trace their current postal addresses. Starting in April 2018, up to five mailings to a large number of Swiss city or village

administrations were needed to obtain as much as possible up-to-date current addresses.

In the spring of 2019, letters providing detailed information about the follow-up study were sent to supposedly correct addresses. Participants who had provided written informed consent for genetic analysis of PBNCs were mailed the postal questionnaire, and those who did not return the questionnaire were mailed a reminder. The last questionnaires were returned in December 2019. Data were coded and anonymously stored in an Excel database.

Ascertainment of diagnosis

The diagnosis AS in the first phase of the study in 1985 was based on the clinical findings and the evaluation of the pelvic radiographs. Probands were categorised according to the mNY criteria, and their FDR were considered to have AS if they met the mNY criteria.

In the follow-up study (2019), recent imaging of the SI joints was mostly unavailable for us to establish whether the FDR who reported having been diagnosed (between 1986 and 2019) by their Swiss rheumatologists to have axSpA met the mNY criteria or had nr-axSpA. Therefore, we considered all new cases in the follow-up study to be suffering from axSpA, that is, they may have either radiographic axSpA or nr-axSpA.

Statistical analysis

Counts were compared by χ^2 testing. The test results are expressed as p values. OR were calculated with 95% CI.

Patient and public involvement

Two patients/coauthors were fully involved in the study.

RESULTS

Altogether 1178 persons, including 363 probands, participated in the first (1985) phase of the study of whom 485 consenting persons could be retrieved for the 2018–2019 follow-up study (125 probands and 360 FDR). Altogether 162 former participants (123 probands and 39 FDR) were known to have died; information about causes of death was not available. Demographic data of the 485 participants who were available and consented to participate in the follow-up study are shown in [table 1](#), together with their radiographic and HLA-B27 status ([figure 1](#)). At baseline (1985) 84/125 (67.2%) probands met mNY criteria, 41 were categorised as nr-axSpA.

Occurrence of axSpA among FDR

The risk to develop axSpA for HLA-B27(+) FDR of HLA-B27(+) probands is considerable. At follow-up (2018–19) 42/162 HLA-B27(+) FDR had been diagnosed as having the disease, that is, an incidence of 25.9%; 95%CI 19.2% to 32.6%. The sex ratio was about equal: 17/67 (25.4%) HLA-B27(+) males and 25/95 (26.3%) HLA-B27(+) females. In contrast, 1/141 (0.7%) of HLA-B27(-) FDR of HLA-B27(+) probands and 1/29 (3.4%) HLA-B27(-) FDR of HLA-B27(-) probands (a son of a nr-axSpA proband) were diagnosed as having axSpA. The first mentioned HLA-B27(-) axSpA case comprises a sister of an HLA-B27(+) proband with nr-axSpA. The HLA-B27 and radiographic sacroiliac status of the proband of a third HLA-B27(-) case (a female) is unknown. Of the 42 HLA-B27(+) cases, 7 had radiographic sacroiliitis in 1985, the remaining 35 either had negative pelvic X-rays or did not undergo radiographic examination of their SI joints at baseline because they were pregnant or <18 years of age at that time. Thus, the diagnosis axSpA of these 35 FDR has

Table 1 Demographic data of axSpA probands and relatives participating in the 2019 Swiss Ankylosing Spondylitis Follow-up Family Study by HLA-B27 status and presence of sacroiliitis by New York criteria*

	No	Males	Females	Mean age (yr) (2019)	Age SD
All participants	485	238	247	64.56	9.73
All axSpA probands	125†‡§	78‡	47§	72.78	7.31
HLA-B27 positive axSpA Probands	110	73	37	72.51	7.16
Sacroiliitis present (1985)	79	54	25		
Sacroiliitis absent	31	19	12		
HLA-B27 negative axSpA probands	13	4	9	75.54	8.68
Sacroiliitis present (1985)	3	2	1		
Sacroiliitis absent	10	2	8		
All relatives	360	160	200	61.70	8.84
All relatives with axial SpA	45	18	27	58.02	8.04
HLA-B27 positive relatives	42	17	25	57.72	8.18
Sacroiliitis present (1985)	7	3	4		
Sacroiliitis absent (1985)	28	12	16		
Sacroiliitis unknown	7	2	5		
HLA-B27 negative relatives	3	1	2	62.0	5.29
Sacroiliitis absent (1985)	3	1	2		
All healthy relatives	315¶	142	173	62.20	8.81
Healthy relatives of HLA-B27 +Probands	262	118	144	62.07	9.17
HLA-B27 positive relatives	120	50**	70††	61.26	8.94
HLA-B27 negative relatives	140	66‡‡	74§§	62.87	9.25
HLA-B27 unknown relatives	3	2¶¶	0	55.50	14.85
Healthy relatives of HLA-B27- probands	29	16	13	60.76	6.66
HLA-B27 positive relatives	0				
HLA-B27 negative relatives	29	16	13	60.76	6.66
Healthy relatives of HLA-B27? probands	24	8	16	65.34	6.00
HLA-B27 positive relatives	9	4	5	68.12	5.69
HLA-B27 negative relatives	15	4	11***	63.67	5.39

New York criteria (grade 2 or higher bilaterally, or grade 3 or 4 unilaterally).

*Sacroiliitis: pelvic radiography from 1985 AS Family Study showing sacroiliitis by modified.

†Altogether 84 (67.2%) probands met modified New York criteria, 41 were categorised as nr-axSpA.

‡Unknown HLA-B27 status of one male proband with sacroiliitis.

§Unknown HLA-B27 status of one female proband with sacroiliitis.

¶HLA-B27 status of the probands of two FDR unknown (one relative HLA-B27(-); one relative HLA-B27(+)).

**No pelvic radiograph available for five males.

††No pelvic radiograph available for 11 females.

‡‡No pelvic radiograph available for four males.

§§No pelvic radiograph available for five females.

¶¶No pelvic radiograph available for one male.

***No pelvic radiograph available for two women.

AS, ankylosing spondylitis; axSpA, axial spondyloarthritis; FDR, first-degree relatives.

been established at some point in time *between* 1986 and 2019. For two of these 35 FDR, recent radiographs were available that confirmed AS by mNY criteria, but for the other 33 FDR no recent imaging of the SI joints were available for us to review at follow-up. Of note, based on the responses to the questionnaire, 38 of all 42 (90.5%) HLA-B27(+) FDR with axSpA fulfilled the ASAS classification criteria for axSpA.^{12 13}

Relationship with radiographic status of proband

In total 37 (88.1%) of the 42 HLA-B27(+) axSpA probands of the 42 HLA-B27(+) affected FDR, met the mNY criteria, not significantly different from the HLA-B27(+) proportion observed among axSpA probands who did not have an affected FDR (210/263 (79.8%), $p=0.20$; OR=1.87, 95% CI (0.70 to 4.98)). Coexistence of both types of the disease (radiographic and non-radiographic) occurred in about 10% of the families.

AAU associated with axSpA

Symptoms suggestive of AAU (one or more episodes of unilateral painful inflamed red eye, and prescription of eye drops containing steroids) occurred significantly more often among FDR who during follow-up developed axSpA than among those who did not: 13/34 (38.2%) vs 29/251 (11.6%), $p=0.00004$; OR=4.74, 95% CI (2.15 to 10.47). Comparing only HLA-B27(+) FDR, 13/34 (38.2%) HLA-B27(+) FDR who developed axSpA had symptoms suggestive of AAU in contrast to 14/114 (12.3%) HLA-B27(+) FDR without axSpA ($p=0.0006$, OR=4.42, 95% CI (1.82 to 10.76)). The prevalence of AAU symptoms among FDR with axSpA (38.2%) is similar to the prevalence of AAU among HLA-B27(+) probands: 35/79 (44.3%) for HLA-B27(+) probands, $p=0.55$). The low prevalence (12.3%) of AAU symptoms in HLA-B27(+) FDR without axSpA does not differ significantly from the 10.9% (15/137) figure for healthy HLA-B27(-) FDR ($p=0.74$).




Figure 1 Flow chart of probands with axial spondyloarthritis (axSpA) by HLA-B27 and radiographic status. First-degree relatives are classified as axSpA or healthy and HLA-B27 status. axSpA, axial spondyloarthritis; HLA-B27+, HLA-B27 positive; HLA-B27-, HLA-B27 negative; HLA-B27?, HLA-B27 unknown; nr, -axSpA, non-radiographic axSpA.

Back pain as a predictor for axSpA

At baseline (1985) CIBP¹⁴ was reported significantly more often (46/286, or 16.1%) by HLA-B27(+) FDR without radiographic sacroiliitis (mean age \pm SD 28.3 \pm 8.1 year) than by 27/272 (9.9%) healthy HLA-B27(-) FDR (mean age \pm SD 28.4 \pm 7.4 year) (OR=1.74, p=0.031). Nonetheless, the onset of CIBP at early age was not a reliable predictor of axSpA. Of the 35 HLA-B27(+) new cases of axSpA who developed the disease between 1986 and 2019 and who had normal SI joints in 1985, 31 had complete data for CIBP at baseline and follow-up. CIBP at baseline occurred in 5/31 (16.1%), not very different from the prevalence of CIBP in 1985 among healthy HLA-B27(+) FDR who never developed the disease (18/126, 14.3%) (OR=1.15, p=0.8). Thus, for FDR of AS probands, the presence of CIBP at young age (mean 28 years) does not reliably predict the long-term development of axSpA.

Sensitivity and specificity of clinical features

The clinical history serves as a useful tool in discriminating people with and without axSpA. Items with sensitivity or specificity above 70% for the presence of axSpA are provided in table 2. It shows that, overall sensitivity decreases as specificity increases, and vice versa. Sensitivity was assessed among all AS probands who met the mNY criteria, whereas specificity was appraised in the group of HLA-B27(-) FDR as they are at very low risk of developing the disease. The best performing items cover three anatomical regions: lumbar spine and gluteal region, thoracic spine, and frontal (anterior) chest wall. These three regions contribute diagnostically about equally (table 2).

The diagnostic yield, in particular specificity, could meaningfully be improved by combining the three anatomical regions into one index of three questions: (1) CIBP; (2) complaints at the thoracic region and (3) complaints at the frontal chest wall. Each of three drawings in the questionnaire pointed to the specific location. Two or three of these questions were answered 'yes' by 69/83 AS probands, providing a sensitivity of 83.1% (figure 2). The specificity among the healthy HLA-B27(-) FDR was 116/133 (87.2%) (OR 33.69, 95% CI 15.6 to 72.5) (p<0.00001).

In the whole group of 45 FDR who had been diagnosed as having axSpA at some point in time between 1985 and 2019, 21/44 (47.7%) FDR fulfilled criteria for CIBP; 28/40 (70.0%) reported thoracic complaints; 21/42 (50%) had complaints at the frontal chest wall. The proportion reporting positive answers to ≥ 2 questions of the three-item composite index among these 'new' axSpA patients was 61.0% (25/41). The specificity of this composite index is 86.3% (101/117) among the HLA-B27(+) healthy FDR.

AxSpA symptoms in 'healthy' relatives

For this analysis, we define 'healthy' to mean never having been diagnosed as suffering from axSpA. Feelings of pain, stiffness or discomfort at the frontal chest wall, the lumbar or thoracic spine occurred in up to 25% of 'healthy' HLA-B27(+) FDR, significantly more than in the HLA-B27(-) healthy relatives (who reported such features in less than 9%) (table 3). These symptoms suggest an inflammatory component for their discomfort. These symptoms tend to cluster in individuals in one anatomical region; in particular complaints at the thoracic spine (last 4 items of table 3) occurred in 9/117 (7.7%) 'healthy' HLA-B27(+) FDR vs only 2/138 (1.4%) of HLA-B27(-) FDR (p=0.015). This suggests possible 'hidden' (undiagnosed) axSpA clinical features, mainly at the thoracic spine and mostly without the well-known CIBP, among genetically predisposed relatives.

DISCUSSION

AS occurs commonly among FDR of HLA-B27(+) probands when they share this susceptibility allele. In this family study spanning 35 years, we observed that \sim 25% of HLA-B27(+) FDR developed axSpA irrespective of their proband's radiographic status (ie, presence or absence of sacroiliitis), but very rarely among the HLA-B27(-) FDR (3% in this study). Moreover, we observed that AS and nr-axSpA may run in one and the same family. This supports the view of genetic homogeneous propensity of both expressions (radiographic and non-radiographic forms) of the disease, at least where the family involved carries HLA-B27. It illustrates a major role of the family history and

Table 2 Features of clinical history discriminating between AS probands meeting modified New York criteria (sensitivity) and healthy HLA-B27 negative relatives (specificity)

Question	AS Probands Sensitivity no/total (%)		HLA-B27 negative Relatives Specificity no/total (%)		OR (95% CI)	P value
Lumbar Spine region	64/87	73.6	117/138	84.8	15.50 (7.97 to 30.16)	<0.00001
Chronic inflammatory low backpain (Calin)						
Insidious onset of back pain	72/86	83.7	80/138	58.0	7.09 (3.65 to 13.79)	<0.00001
More than 3 months duration	67/86	77.9	103/138	74.6	10.38 (5.48 to 19.63)	<0.00001
Associated with morning stiffness >30 min	66/86	76.7	102/138	73.9	9.35 (4.99 to 17.52)	<0.00001
Back pain starting before age 40	84/86	97.7	88/138	63.8	73.92 (17.43 to 33.45)	<0.00001
Pain and stiffness	68/87	78.2	105/138	76.1	11.39 (5.99 to 21.63)	<0.00001
Worsening at early morning	56/86	65.1	125/137	91.2	19.44 (9.28 to 40.75)	<0.00001
Waking up during the night because of complaints	56/86	65.1	122/138	88.4	14.23 (7.18 to 28.22)	<0.00001
Leaving bed and walking around during the night	50/84	59.5	131/136	96.3	38.53 (14.26 to 104.08)	<0.00001
Pain irradiating into gluteal region	64/82	78.0	91/138	65.9	6.88 (3.66 to 12.93)	<0.00001
Usage of analgesics	82/86	95.3	86/138	62.3	33.90 (11.73 to 97.96)	<0.00001
Relief of symptoms due to analgesics	73/88	83.0	88/137	64.2	8.74 (4.53 to 16.85)	<0.00001
Thoracic region						
Complaints at thoracic region ever	67/84	79.8	106/134	79.1	14.92 (7.59 to 29.33)	<0.00001
Pain and stiffness	47/88	53.4	131/138	94.9	21.45 (9.01 to 51.11)	<0.00001
Complaints at early morning	54/86	62.8	116/138	84.1	8.90 (4.73 to 16.73)	<0.00001
Morning stiffness at thoracic spine region	45/84	53.6	132/137	96.4	30.46 (11.31 to 82.03)	<0.00001
Complaints if body position does not change	52/83	62.7	119/137	86.9	11.09 (5.70 to 21.58)	<0.00001
Usage of analgesics	48/85	56.5	129/137	94.2	20.92 (9.09 to 48.12)	<0.00001
Relief of symptoms due to analgesics	46/87	52.9	130/138	94.2	18.23 (7.96 to 41.76)	<0.00001
Ventral chest wall region						
Complaints at ventral chest wall region ever	62/86	72.1	113/136	83.1	12.69 (6.62 to 24.32)	<0.00001

Included items are presented by anatomical region and have sensitivity and/or specificity >70%.
AS, ankylosing spondylitis.

putative additional familial factors as predictor of development of axSpA.^{5 15 16}

One might wonder whether the cumulative axSpA incidence of ~25% for HLA-B27(+) FDR might be inflated by *selection bias* or case-ascertainment (clinical examination at baseline and patient reported information at follow-up).




Figure 2 The clinical history can be used as a diagnostic tool. Sensitivity and specificity of complaints at three anatomical regions are provided for each region separately and for a composite index of complaints at these three regions (right). Complaints of the lumbar spine were assessed by chronic inflammatory back pain (positive if ≥ 4 (of 5) Calin items are met). The thoracic spine was evaluated by complaints of pain and discomfort at the dorsal spine. Pain and discomfort at the ventral chest wall were likewise appraised. The three region index was considered positive if complaints at ≥ 2 regions were present.

Selection bias would have occurred if symptomatic HLA-B27(+) FDR who had developed axSpA after baseline would have preferentially volunteered to participate in the follow-up study. In fact, the proportion participating HLA-B27(+) FDR was even slightly lower compared with baseline: in 1985 360/668 (53.9%) FDR of HLA-B27(+) probands were HLA-B27(+) compared with 162/329 (49.2%) at follow-up.

Ascertainment of diagnosis at baseline (clinical diagnosis at the study centre) differed from ascertainment at follow-up (patient reported diagnosis by a Swiss rheumatologist). Moreover, the concept of disease has broadened in the last two decades. This may have impacted (inflated) our axSpA incidence ~25% figure. However, as our findings illustrate, 'AS without sacroiliitis' was already known in 1985,⁸ and the Rome criteria¹⁷ allow the diagnosis if 4 of 5 clinical criteria are met in the absence of definite sacroiliitis. In recent decades the broader concept of axSpA has become widely accepted. The increased awareness and improved imaging (MRI) have promoted better recognition of axSpA. Taking the differences in case-ascertainment into account, a somewhat higher prevalence figure than the earlier reported 21% figure⁵ seems sensible.

The occurrence at baseline of 'AS without radiographic sacroiliitis' among axSpA probands, shown in table 1, illustrates that this entity is nowadays widely known as nr-axSpA.

Uveitis as a predictor for axSpA

AAU is a common, usually unilateral, HLA-B27 associated, intraocular inflammatory disease, concomitantly occurring in patients with axSpA. The prevalence of AAU in AS increases with disease duration, and may exceed 50%, particularly in HLA-B27(+) patients with longstanding disease.¹⁸⁻²³ We accepted

Table 3 Prevalence of symptoms suggesting 'hidden' axial SpA among HLA-B27 positive FDR compared with HLA-B27 negative FDR

Question	Relatives HLA-B27 positive no positive/total and percentage		Relatives HLA-B27 negative no positive/total and percentage		OR	95% CI	P value
Change of body position causes exacerbation of complaints at the ventral chest wall	6/114	5.3	1/138	0.7	7.61	0.90 to 64.18	0.029
Worsening of low back pain during the early morning before getting up	28/111	25.2	12/137	8.8	3.51	1.69 to 7.30	0.00046
Persisting low back pain after getting up	19/117	16.2	11/137	8.0	2.22	1.01 to 4.88	0.043
Waking up regularly after 14:00 hour because of pain or discomfort at the thoracic spine	7/116	6.0	1/137	0.7	8.73	1.06 to 72.07	0.016
Morning stiffness at the thoracic spinal region	11/117	9.4	5/137	3.6	2.74	0.92 to 8.13	0.06
Morning stiffness at the thoracic spinal region for at least 15 to 60 min	8/117	6.8	2/138	1.4	4.99	1.04 to 23.99	0.027
Improvement of stiffness at the thoracic spinal region through exercises	12/112	10.7	5/134	3.7	3.10	1.06 to 9.08	0.032
Exercising improves thoracic complaints due to sustained body position	17/116	14.7	9/136	6.6	2.42	1.04 to 5.67	0.037

None of these FDR of HLA-B27 positive axSpA probands had ever been diagnosed as axSpA. axSpA, axial spondyloarthritis; FDR, first-degree relatives.

participants' reply of having used corticosteroid containing eye drops as the most important proxy for a diagnosis of AAU. Although this might have face validity, it is not a firm proof. With this limitation in mind, AAU was significantly associated with development of axSpA, whereas there was no difference in occurrence of AAU between healthy HLA-B27(+) and HLA-B27(-) FDR. This is compatible with the reported 45% association of HLA-B27 with AAU in the general population, that is, about equally in HLA-B27(+) and HLA-B27(-) individuals.²⁴

The stronger association of AAU with axSpA than with HLA-B27 might, at least partly, be due to diagnostic suspicion bias, that is, an established diagnosis (say axSpA) might increase the likelihood that an associated disease (say AAU) will be diagnosed appropriately. Nonetheless, our findings are strongly supported by recent genome-wide association studies revealing that the association of AAU and axSpA is primarily with the full genetic overlap (HLA and non-MHC genes) of both conditions rather than with the HLA-B27 allele alone.^{15,23} One may conclude that AAU is more closely related to the rheumatologic disease than to the HLA-B27 allele, and therefore, may predict future axSpA. Literature indicates that among HLA-B27(+) AAU patients the prevalence of concomitant axSpA may rise to about 80%, illustrating that occurrence of AAU among relatives warrants screening for axSpA.¹⁹

Chronic back pain as a predictor for axSpA

CIBP at young age failed to predict future development of axSpA in our study. Reveille *et al*²⁵ reported that FDR of AS patients had earlier onset of CIBP than those with non-IBP, and that their CIBP persisted longer (73% of the FDR still had CIBP after a mean of 5.5 years). On the other hand, Wang *et al*²⁶ found that a minority of patients with new onset IBP progressed to SpA, while IBP resolved in many. They suggest that the finding that IBP often resolves may explain the difference between the prevalence of IBP (3%–6%) and the prevalence of SpA (0.4%–1.3%). Of note, at baseline 16.1% (46/268) of our HLA-B27(+) FDR had CIBP without sacroiliitis vs 9.9% (27/272) of the HLA-B27(-) FDR ($p=0.03$), but there was no difference observed at follow-up 35 years later. This supports the view that CIBP might be a potentially resolving manifestation of axSpA, that is, a favorable outcome of the disease. This view is also supported by recently published findings from the pre-SpA cohort study. IBP was reported by 19% of seemingly healthy FDR (of HLA-B27(+)

axSpA patients) with an equal distribution between HLA-B27(+) and HLA-B27(-) FDR. Progression from subclinical inflammation to clinical axSpA within 1 year of follow-up occurred in 6%, and was mainly (86%) observed in HLA-B27(+) FDR.²⁰

Clinical history as an aid to diagnosis

For axSpA the diagnostic yield and accuracy might improve by paying close attention not only to the well-known inflammatory symptoms at the lumbar spine,^{14,27} but also to complaints of pain and discomfort at the thoracic spinal and the frontal chest wall region. Combining features at these three anatomical regions into a *triad* yielded high (87%) specificity. While our study requires independent validation, this finding compares quite well with Rudwaleit's refined set of criteria for CIBP that have a sensitivity of 70.3% and specificity of 81.2% if at least two of the four parameters (morning stiffness of >30 min' duration, age at onset of back pain, no improvement in back pain with rest, awakening because of back pain during the second half of the night only, alternating buttock pain and time period of the onset of back pain) were fulfilled.²⁷ It would be worthwhile to combine and evaluate the refined CIBP criteria with complaints of pain and discomfort at the thoracic spine and the chest wall. If such studies would also demonstrate high specificity and sufficient sensitivity, then the composite triad index of complaints of pain and discomfort at lumbar spine, thoracic spine and chest wall might also provide the needed improvement of the specificity of the current classification criteria for axSpA that is being addressed in the ongoing CLASSIC study.^{28,29}

In summary, about 25% of FDR of HLA-B27(+) probands with axSpA also develop the disease. Occurrence of AAU among FDR of patients with axSpA calls for screening for the disease, but presence of CIBP among young FDR has no long-term predictive value for the diagnosis axSpA. Paying attention to prevailing symptoms of pain and discomfort at the thoracic spine and the frontal chest wall in addition to CIBP may improve the diagnostic yield and classification of axSpA. 'Healthy' HLA-B27(+) FDR with such symptoms may have unnoticed 'hidden' axSpA, that may not be captured by the current sets of criteria or fall short of being diagnosed properly.

Author affiliations

¹University of Bern, Bern, Switzerland

²Department of Internal Medicine, University of Maastricht, Maastricht, The Netherlands

³Case Western Reserve University School of Medicine, Cleveland, Ohio, USA

⁴Faculty of Health, School of Biomedical Sciences, Queensland University of Technology, Brisbane, Queensland, Australia

⁵Centre for Genomics and Personalised Health, Queensland University of Technology, Brisbane, Queensland, Australia

⁶SVMB, Flims, Switzerland

⁷Rheumatology Research, Mortroux, Belgium

⁸Department of Mathematical Sciences, Kent State University, Kent, Kent, USA

⁹Department of Rheumatology and Clinical Immunology, Medical Center Monbijou, Meikirch, Switzerland

¹⁰Department of Medical and Molecular Genetics, King's College London, London, UK

¹¹Genomics England, London, UK

Correction notice This article has been corrected since it published Online First. Figure 1 has been replaced and affiliation 10 updated.

Acknowledgements We like to thank all patients, spouses and relatives for their kind cooperation and are grateful to the Swiss city and village administrations for retrieving current addresses. We also thank Hans-Ueli Rentsch*, MD, Hans Valkenburg*, MD, Arnold Cats*, MD, Herman Kroon, MD and Niklaus Gerber, MD for their contributions in performing the study. Caroline Kaegi provided helpful secretarial assistance. (*Deceased)

Contributors All authors had full access to all of the data in the study and take responsibility for the data and the accuracy of the data analysis. Concept and design: SMvdL, MAK, HB, HvZ and MAB conceived the study. SMvdL, HB and HvZ designed the questionnaire. Drafting of the manuscript: SMvdL, MAK and MAB. Genetical analysis: ZL and MAB. Statistical analysis: SMvdL, ZL, KK and MAB. Obtained funding: SMvdL and MAB. Administrative, technical or material support: HvZ and PMV. Acquisition of data: SMvdL, HvZ and PMV. Critical revision of the manuscript for important intellectual content: all authors. Guarantor: SMvdL.

Funding The 1985 baseline study was funded by the Swiss National Fund, Schweizer Rück Insurance and Ciba-Geigy, Switzerland. The 2018 follow-up study was funded/supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London and/or the NIHR Clinical Research Facility.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by Ethical Committee of the Swiss Kanton Bern (2017-00536). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Not applicable.

ORCID iDs

Sjef M van der Linden <http://orcid.org/0000-0002-0087-0351>

Muhammad Asim Khan <http://orcid.org/0000-0003-4704-8311>

Zhixiu Li <http://orcid.org/0000-0002-2924-9120>

Peter M Villiger <http://orcid.org/0000-0002-8859-9964>

Matthew A Brown <http://orcid.org/0000-0003-0538-8211>

REFERENCES

- de Blecourt J, Polman A, de Blecourt-Meindersma. Hereditary factors in rheumatoid arthritis and ankylosing spondylitis. *Ann Rheum Dis* 1961;20:215–23.
- Carter N, Williamson L, Kennedy LG, et al. Susceptibility to ankylosing spondylitis. *Rheumatology* 2000;39:445.
- Brown MA, Laval SH, Brophy S, et al. Recurrence risk modelling of the genetic susceptibility to ankylosing spondylitis. *Ann Rheum Dis* 2000;59:883–6.
- Calin A, Marder A, Becks E, et al. Genetic differences between B27 positive patients with ankylosing spondylitis and B27 positive healthy controls. *Arthritis Rheum* 1983;26:1460–4.
- van der Linden SM, Valkenburg HA, de Jongh BM, et al. The risk of developing ankylosing spondylitis in HLA-B27 positive individuals. A comparison of relatives of spondylitis patients with the general population. *Arthritis Rheum* 1984;27:241–9.
- Robinson WP, van der Linden SM, Khan MA, et al. HLA-Bw60 increases susceptibility to ankylosing spondylitis in HLA-B27+ patients. *Arthritis Rheum* 1989;32:1135–41.
- Wellcome Trust Case Control Consortium, Australo-Anglo-American Spondylitis Consortium (TASC), Burton PR, et al. Association scan of 14,500 nonsynonymous SNPs in four diseases identifies autoimmunity variants. *Nat Genet* 2007;39:1329–37.
- Khan MA, van der Linden SM, Kushner I, et al. Spondylitic disease without radiologic evidence of sacroiliitis in relatives of HLA-B27 positive ankylosing spondylitis patients. *Arthritis Rheum* 1985;28:40–3.
- Rudwaleit M, Khan MA, Sieper J. The challenge of diagnosis and classification in early ankylosing spondylitis: do we need new criteria? *Arthritis Rheum* 2005;52:1000–8.
- van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the new York criteria. *Arthritis Rheum* 1984;27:361–8.
- van Hoveen L, Luime J, Han H, et al. Identifying axial spondyloarthritis in Dutch primary care patients, ages 20-45 years, with chronic low back pain. *Arthritis Care Res* 2014;66:446–53.
- Rudwaleit M, Landewé R, van der Heijde D, et al. The development of assessment of spondyloarthritis International Society classification criteria for axial spondyloarthritis (part I): classification of paper patients by expert opinion including uncertainty appraisal. *Ann Rheum Dis* 2009;68:770–6.
- Rudwaleit M, van der Heijde D, Landewé R, et al. The development of assessment of spondyloarthritis International Society classification criteria for axial spondyloarthritis (part II): validation and final selection. *Ann Rheum Dis* 2009;68:777–83.
- Calin A, Porta J, Fries JF, et al. Clinical history as a screening test for ankylosing spondylitis. *JAMA* 1977;237:2613–4.
- Huang X-F, Li Z, De Guzman E, et al. Genomewide association study of acute anterior uveitis identifies new susceptibility loci. *Invest Ophthalmol Vis Sci* 2020;61:3.
- van Lunteren M, Sepriano A, Landewé R, et al. Do ethnicity, degree of family relationship, and the spondyloarthritis subtype in affected relatives influence the association between a positive family history for spondyloarthritis and HLA-B27 carriage? Results from the worldwide ASAS cohort. *Arthritis Res Ther* 2018;20:166.
- Kellgren J, Jeffrey M, Ball J. *The epidemiology of chronic rheumatism*. Blackwell: Oxford, 1963.
- Khan MA, Kushner I, Braun WE. Comparison of clinical features in HLA-B27 positive and negative patients with ankylosing spondylitis. *Arthritis Rheum* 1977;20:909–12.
- Juanola X, Loza Santamaria E, Cordero-Coma M, et al. Description and prevalence of spondyloarthritis in patients with anterior uveitis: the sentinel interdisciplinary collaborative project. *Ophthalmology* 2016;123:1632–6.
- de Jong HMY, de Winter JH, van der Horst-Bruinsma IE, et al. Progression from subclinical inflammation to overt SpA in first degree relatives of spa patients is associated with HLA-B27: the Pre-SpA cohort. *Arthritis Care Res* 2021. doi:10.1002/acr.24743. [Epub ahead of print: 05 Jul 2021].
- Sharma SM, Jackson D. Uveitis and spondyloarthropathies. *Best Pract Res Clin Rheumatol* 2017;31:846–62.
- Khan MA, Haroon M, Rosenbaum JT. Acute anterior uveitis and spondyloarthritis: more than meets the eye. *Curr Rheumatol Rep* 2015;17:59.
- Robinson PC, Clauhsuis TAM, Cortes A, et al. Genetic dissection of acute anterior uveitis reveals similarities and differences in associations observed with ankylosing spondylitis. *Arthritis Rheumatol* 2015;67:140–51.
- Beckingsale AB, Davies J, Gibson JM, et al. Acute anterior uveitis, ankylosing spondylitis, back pain, and HLA-B27. *Br J Ophthalmol* 1984;68:741–5.
- Doughem K, Weisman MH, Ward MM, et al. Chronic back pain in first-degree relatives (FDRs) of patients with ankylosing spondylitis: predictive value of HLA-B27 and persistence of inflammatory back pain over time. *RMD Open* 2020;6:e001418.
- Wang R, Crowson CS, Wright K, et al. Clinical evolution in patients with new-onset inflammatory back pain: a population-based cohort study. *Arthritis Rheumatol* 2018;70:1049–55.
- Rudwaleit M, Metter A, Listing J, et al. Inflammatory back pain in ankylosing spondylitis: a reassessment of the clinical history for application as classification and diagnostic criteria. *Arthritis Rheum* 2006;54:569–78.
- Khan MA, van der Linden S. Axial spondyloarthritis: a better name for an old disease: a step toward uniform reporting. *ACR Open Rheumatol* 2019;1:336–9.
- van der Linden S, Akkoc N, Brown MA, et al. The ASAS criteria for axial spondyloarthritis: strengths, weaknesses, and proposals for a way forward. *Curr Rheumatol Rep* 2015;17:62.